

REPORT DOCUMENTATION PAGE					Form Approved OMB No. 0704-0188	
<small>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to the Department of Defense, Executive Services and Communications Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</small> PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION.						
1. REPORT DATE (DD-MM-YYYY) 15-02-2012		2. REPORT TYPE Conference Proceedings		3. DATES COVERED (From - To)		
4. TITLE AND SUBTITLE Characterization of Suspended Particulates in the Northern Gulf of Mexico from Ocean Color Remote Sensing				5a. CONTRACT NUMBER		
				5b. GRANT NUMBER		
				5c. PROGRAM ELEMENT NUMBER 0601153N		
6. AUTHOR(S) ZhongPing Lee, C. Huang, B. Lubac, L. Guo, Dong Shan Ko, S. Lohrenz, R. Gould				5d. PROJECT NUMBER		
				5e. TASK NUMBER		
				5f. WORK UNIT NUMBER 73-9857-00-5		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory Oceanography Division Stennis Space Center, MS 39529-5004				8. PERFORMING ORGANIZATION REPORT NUMBER NRL/PP/7320-10-0512		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research One Liberty Center 875 North Randolph Street, Suite 1425 Arlington, VA 22203-1995				10. SPONSOR/MONITOR'S ACRONYM(S) ONR		
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution is unlimited.						
13. SUPPLEMENTARY NOTES 20120217344						
14. ABSTRACT The ecosystem in the Northern Gulf of Mexico (NGOM) is composed of fresh waters from coastal run-offs and the Mississippi River as well as clear-salty waters from the gulf. As a result, the composition of water constituents, as well as their spatial and temporal distributions, is complex and heterogeneous, with severe hypoxia events constantly happening in the NGOM region. In this study, using properties derived from measurements of ocean color radiance by a satellite sensor, the nature of the suspended particulates was characterized. In particular, not only the concentrations (and their spatial distributions) of suspended particulate matters were derived from satellite images, but also the organic/inorganic nature of these particulates were characterized with derived inherent optical properties. These results could help us better understand the dynamics of coastal ecosystems and the development of hypoxia.						
15. SUBJECT TERMS coastal ecosystem, hypoxia, satellite measurement, suspended particulates						
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 7	19a. NAME OF RESPONSIBLE PERSON Dong Shan Ko	
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NUMBER (Include area code) 228-688-5448	

Characterization of Suspended Particulates in the Northern Gulf of Mexico from Ocean Color Remote Sensing

Zhongping Lee, GRI, Mississippi State University, Stennis Space Center, MS 39529
Changchun Huang, GRI, Mississippi State University, Stennis Space Center, MS 39529
Bertrand Lubac, GRI, Mississippi State University, Stennis Space Center, MS 39529
Laodong Guo, DMS, University of Southern Mississippi, Stennis Space Center, MS 39529
Dongshan Ko, Naval Research Laboratory, Stennis Space Center, MS 39529
Steven Lohrenz, DMS, University of Southern Mississippi, Stennis Space Center, MS 39529
Richard Gould, Naval Research Laboratory, Stennis Space Center, MS 39529

ABSTRACT

The ecosystem in the Northern Gulf of Mexico (NGOM) is composed of fresh waters from coastal run-offs and the Mississippi River as well as clear-salty waters from the gulf. As a result, the composition of water constituents, as well as their spatial and temporal distributions, is complex and heterogeneous, with severe hypoxia events constantly happening in the NGOM region. In this study, using properties derived from measurements of ocean color radiance by a satellite sensor, the nature of the suspended particulates was characterized. In particular, not only the concentrations (and their spatial distributions) of suspended particulate matters were derived from satellite images, but also the organic/inorganic nature of these particulates were characterized with derived inherent optical properties. These results could help us better understand the dynamics of coastal ecosystems and the development of hypoxia.

INTRODUCTION

The northern Gulf of Mexico is not only economically important, but also dynamic in ecosystem structures. It receives freshwater discharges and material export through Mobile Bay estuarine and river plumes from the Mississippi River and Atchafalaya River, the largest river system in the North America. The interactions between these river discharges and the relatively clear open gulf waters nurture the largest seafood industry [Ache *et al.*, 2008] in the United States, and form one of the most severe hypoxia region [Rabalais *et al.*, 2002b; Walker and Rabalais, 2006]. Indeed, the NGOM ecosystem is sensitive to environmental and climate changes [Burkart and James, 1999; Rabalais *et al.*, 2002a; Raymond *et al.*, 2008], and is experiencing profound influences by nutrients and particulate matter loadings from river runoff and coastal erosion, causing hypoxia, eutrophication, other potential environmental problems [Brunner *et al.*, 2006; Rabalais *et al.*, 2002a], and the enormous impacts from the Deep Water Horizon tragedy. On the other hand, because of the complex composition of various constituents, NGOM waters also post enormous challenges to propel the advancement of ocean color remote sensing. Any techniques/algorithms developed and proved working with NGOM waters will have significant impact on ocean color remote sensing of global coastal waters.

One important parameter for ecosystem studies is the concentration of suspended particulate matters, as it not only affects subsurface light field (then photosynthesis), it could also be a tracer of pollutant transport and an indicator of shoreline erosion. Adequate information of the amount, nature, and spatial variation of suspended sediments is highly desired for marine biology and geology, and for coastal management. Ship survey generally provides detailed information of discrete sampling locations. But for broad coastal areas, remote sensing by satellite sensors is the only feasible means to obtain synoptic information of the distribution of suspended sediments.

Traditional remote sensing studies of suspended particulates, however, have been focused on the estimation of total suspended sediments (or total suspended solids; practically particles collected on a filter pad) from remotely measured water reflectance [D'Sa *et al.*, 2007a; Dekker *et al.*, 2002; Stumpf, 1988]. This satellite product, however, does not separate mineral sediments from biogenic materials (such as phytoplankton), consequently little is known about the quality of such suspended particulate matters in coastal ecosystems. Note that suspended particulate organic and inorganic matters have different residence time, turnover rates and pathogens, thus contribute differently in carbon cycling, nutrient and phytoplankton bloom dynamics, and toxic metal transport in coastal ecosystems [Baskaran *et al.*, 1996; Buesseler *et al.*, 1992; Dagg *et al.*, 2008; Sridhar *et al.*, 2008; Trefry *et al.*, 1994; Wang *et al.*, 2004]. Therefore, it is not only desired but also important to measure both the loading of suspended particulate matters and the quality (or class) of such materials [Balch *et al.*, 2005; Gould and Arnone, 2003; Stramski *et al.*, 2008].

Here, with measurements made by MERIS over the NGOM waters, we present products that measure the concentration of SPM and the nature (characteristics) of such particulates, and briefly discuss their interactions with currents in coastal regions.



Fig. 1. Northern Gulf of Mexico.




Fig. 2. Spatial distribution of [SPM]. Derived from MERIS (Feb. 6, 2007) $Rrs(665)$ and $Rrs(560)$.

CONCENTRATIONS AND CHARACTERISTICS OF SUSPENDED PARTICULATES FROM REMOTE SENSING

For NGOM waters, Fig. 1 shows a true color image of MERIS measurements (~ 300 m spatial resolution) of Feb. 6, 2007, with Level 1 (calibrated top-of-atmosphere radiance) data obtained from ESA. Remote sensing reflectance (Rrs , defined as the ratio of water-leaving radiance to downwelling irradiance just above the surface) was derived by applying the Case-2 atmosphere correction algorithm developed by Doerffer and Schiller [2006].

In the past decade, various algorithms [Bukata *et al.*, 1995; D'Sa *et al.*, 2007b; Dekker *et al.*, 2002; Miller and McKee, 2004; Stumpf and Pennock, 1989] have been developed to retrieve the loading of the suspended particulate matters ([SPM], units: mg/m^3) in surface waters. Fig. 2 displays the spatial distribution of [SPM] derived with the algorithm of D'Sa *et al.* [2007a], which uses Rrs at 670 nm and 555 nm as inputs (replaced as 665 nm and 560 nm for MERIS measurements) and was specifically tuned for the waters in the northern Gulf of Mexico. The image suggests that except the lower right

corner where impacts from cirrus clouds caused falsely enhanced [SPM], higher concentrations of [SPM] are found, as expected, in coastal regions that are adjacent to land. In addition, eddies resulted from mixing of coastal currents are highlighted in the [SPM] image.

This [SPM] image, however, although shows a decrease gradient of SPM loading from nearshore to offshore, does not reveal particle features of the distinctive water patch (light blue) in the lower left corner. This patch of waters has significantly different spectral R_{rs} than that of adjacent waters with similar R_{rs} values at 665 nm (see Fig. 3). The contrast in spectral R_{rs} indicates that the water constituents of the two pixels (or patches) could be significantly different.




Fig. 3. Examples of spectral R_{rs} .

To evaluate and understand this contrast, at least to the first order, inherent optical properties (IOPs) of waters shown in Fig.1 were derived by applying the Quasi-Analytical Algorithm (QAA) [Lee *et al.*, 2002], with derived total absorption and backscattering coefficients shown in Fig. 4. Apparently, while the absorption image has some resemblance of the [SPM] spatial pattern, the backscattering image show enhanced scattering of the light-blue patch. This contrast in IOPs further indicates different characteristics of suspended particulates in the northern Gulf of Mexico.




Fig. 4. Spatial distribution of absorption coefficient (left) and backscattering coefficient (right).

The difference in particulate characteristics is further manifested with an analysis of the backscattering coefficients of phytoplankton (b_{b-Phy}) and non-phytoplankton (b_{b-Nphy}) components, both were calculated at 560 nm. b_{b-Phy} is based on QAA-derived phytoplankton absorption coefficient at 442 nm ($a_{Phyt}(442)$) with the following steps:

- 1) Index of phytoplankton amount ($[Phyt]$) is estimated based on the absorption model of Bricaud et al [1995]

$$[Phyt] = \left(\frac{a_{phyt}(442)}{0.065} \right)^{1/0.728} \quad (1)$$

- 2) Scattering coefficient (at 560 nm) of phytoplankton is estimated based on the model of Gordon and Morel [1983] with the above $[Phyt]$ as input

$$b_{phyt}(560) = 0.3([Phyt])^{0.62} \quad (2)$$

- 3) Backscattering coefficient (at 560 nm) of phytoplankton $b_{b-phyt}(560)$ is estimated as 1% of $b_{phyt}(560)$.

Further, backscattering coefficient (at 560 nm) of Non-Phytoplankton ($b_{b-NPhyt}(560)$) is derived from QAA derived particle backscattering coefficient ($b_{bp}(560)$),

$$b_{b-NPhyt}(560) = b_{bp}(560) - b_{b-phyt}(560), \quad (3)$$

and a b_b -ratio is evaluated as $b_{b-NPhyt}(560)/b_{b-phyt}(560)$, with results shown in Fig. 5. It is found that generally b_b -ratio is in a range of 2 - 30 for the study area, and higher values (> 50) were found around the Mississippi River delta. Stavn and Riecheter [2008] found a ratio of 4-30 between the scattering coefficients of non-organic to organic particulates for waters around the Mobile Bay. The b_b -ratio results are consistent with those findings.




Fig. 5. Spatial distribution of b_b -ratio.




Fig. 6. NCOM results of coastal currents.

Combining the spatial distribution of $a(442)$, $b_b(442)$ and b_b -ratio, and comparing these patterns with that shown by the $[SPM]$, more information about the characteristics, and spatial variation, emerge. 1) Non-phytoplankton scatterings, mainly dominated by mineral particles, make strong contributions in the coastal regions, especially the waters off Mobile Bay (Alabama). 2) Extremely strong non-phytoplankton scattering appeared in the Mississippi River plume and the Atchafalaya Bay, due likely to resuspended mineral sediments. 3) Waters in the Lake Pontchartrain show suspended particles with

both biogenic and mineral characteristics. And, 4) The light-blue patch in the lower left corner of Fig. 1 is dominated by particles of non-phytoplankton (or mineral) character. This detached mineral-dominant water patch could be resulted from mixings between coastal runoff and ocean currents. During winter time, coastal currents flow westward (see Fig. 6), which could effectively separate the coastal runoff water with ocean water, of which the contribution from non-phytoplankton is much less (see Fig. 5). During the mixing of runoff waters with clearer oceanic waters, likely a dilution effect, or removal of organic coatings of the particles, resulted in a reduction of the absorption coefficient (at 442 nm) in the light-blue patch, consequently enhanced Rrs in the blue-green wavelengths are observed (Fig. 3). At the same time, those particles may add nutrients in the deeper water column and eventually contribute to the occurrence of hypoxia in this region [Rabalais *et al.*, 2002a].

SUMMARY

As an example, concentration of suspended particle matters ([SPM]), along with optical properties, in the northern Gulf of Mexico were derived from a measurement of MERIS (Feb. 6, 2007). Presently [SPM] image products provide valuable spatial variations of particulates in the upper water column. Its nature of derivation, however, could not inform the characteristics of the suspended particulates. With optical properties derived from advanced near-analytical algorithms, such characteristic information could then be evaluated. Combined with circulation models we will be able to better understand and predict the dynamics of particulates and coastal ecosystems.

ACKNOWLEDGEMENT

Supports from NASA's Water- and Energy Cycle Program and from the Northern Gulf Institute are greatly appreciated.

REFERENCE

- Ache, B., D. Bylsma, K. Crossett, C. David, A. Edwards, and N. Wallace (2008), Gulf of Mexico at a Glance., National Ocean Service/NOAA/U.S. Department of Commerce, Washington, D.C.
- Balch, W. M., H. R. Gordon, B. C. Bowler, D. T. Drapeau, and E. S. Booth (2005), Calcium carbonate measurements in the surface global ocean based on Moderate-Resolution Imaging Spectroradiometer data, *J. Geophys. Res.*, *110*(C07001), doi:10.1029/2004JC002560.
- Baskaran, M., P. H. Santschi, L. Guo, T. S. Bianchi, and C. Lambert (1996), ^{234}Th - ^{238}U disequilibria in the Gulf of Mexico: the importance of organic matter and particle concentration, *Cont. Shelf Res.*, *16*, 353-380.
- Bricaud, A., M. Babin, A. Morel, and H. Claustre (1995), Variability in the chlorophyll-specific absorption coefficients of natural phytoplankton: Analysis and parameterization, *J. Geophys. Res.*, *100*, 13321-13332.
- Brunner, C. A., J. M. Beall, S. J. Bentley, and Y. Furukawa (2006), Hypoxia hotspots in the Mississippi Bight, *J. Foraminiferal Res.*, *36*, 95-107.
- Buesseler, K. O., M. P. Bacon, J. K. Cochran, and H. D. Livingston (1992), Carbon and nitrogen export during the JGOFS North Atlantic Bloom Experiment estimated from ^{234}Th - ^{238}U disequilibria, *Deep-Sea Res.*, *39*, 1115-1137.
- Bukata, R. P., J. H. Jerome, K. Y. Kondratyev, and D. V. Pozdnyakov (1995), *Optical Properties and Remote Sensing of Inland and Coastal Waters*, CRC Press, Boca Raton, FL.
- Burkart, M. R., and D. E. James (1999), Agricultural-Nitrogen Contributions to Hypoxia in the Gulf of Mexico, *J. Environ. Qual.*, *28*, 850-859.

- D'Sa, E. J., R. L. Miller, and B. A. McKee (2007a), Suspended particulate matter dynamics in coastal waters from ocean color: Applications to the Northern Gulf of Mexico, *Geophysical Research Letters*, *34*, doi:10.1029/2007GL031192.
- D'Sa, E. J., R. L. Miller, and B. A. McKee (2007b), Suspended particulate matter dynamics in coastal waters from ocean color: Applications to the Northern Gulf of Mexico, *Geophysical Research Letters*, *34*(L23611), doi:10.1029/2007GL031192.
- Dagg, M. J., T. Bianchi, B. McKee, and R. Powell (2008), Fates of dissolved and particulate materials from the Mississippi river immediately after discharge into the northern Gulf of Mexico, USA, during a period of low wind stress, *Continental Shelf Research*, *28*, 1443–1450.
- Dekker, A. G., R. J. Vos, and S. W. M. Peters (2002), Analytical algorithms for lake water TSM estimation for retrospective analyses of TM and SPOT sensor data, *Int. J. Remote Sensing*, *23*(1), 15–35.
- Doerffer, R., H. Schiller, and M. Peters (2006), MERIS Case-2 Regional Processor, edited, pp. BEAM Plug-ins, ESA.
- Gordon, H. R., and A. Morel (1983), *Remote assessment of ocean color for interpretation of satellite visible imagery: A review*, 44 pp., Springer-Verlag, New York.
- Gould, R. W., and R. A. Arnone (2003), Coastal transport of organic and inorganic matter, *2003 NRL Review Book, May 2003*, 199–202.
- Lee, Z. P., K. L. Carder, and R. Arnone (2002), Deriving inherent optical properties from water color: A multi-band quasi-analytical algorithm for optically deep waters, *Applied Optics*, *41*, 5755–5772.
- Miller, R. L., and B. A. McKee (2004), Using MODIS 250 m imagery to map concentrations of total suspended matter in coastal waters, *Remote Sens. Environ.*, *93*, 259–266.
- Rabalais, N. N., R. E. Turner, Q. Dorteh, D. Justic, V. J. B. Jr., and W. J. W. Jr. (2002a), Nutrient-enhanced productivity in the northern Gulf of Mexico: past, present and future, *Hydrobiologia*, *475/476*, 39–63.
- Rabalais, N. N., R. E. Turner, and D. Scavia (2002b), Beyond science into policy: Gulf of Mexico hypoxia and the Mississippi River, *BioScience*, *52*(129–142).
- Raymond, P. A., N.-H. Oh, R. E. Turner, and W. Broussard (2008), Anthropogenically enhanced fluxes of water and carbon from the Mississippi River, *Nature*, *451* 449–452.
- Sridhar, P. N., I. V. Ramana, M. M. Ali, and B. Veerananarayana (2008), Understanding the suspended sediment dynamics in the coastal waters of the Bay of Bengal using high resolution ocean colour data, *CURRENT SCIENCE, VOL. , NO. , 10 JUNE 1501*, *94*(11), 1499–1502.
- Stavn, R. H., and S. J. Richter (2008), Biogeo-optics: particle optical properties and the partitioning of the spectral scattering coefficient of ocean waters, *Appl. Opt.*, *47*, 2660–2679.
- Stramski, D., R. A. Reynolds, M. Babin, S. Kaezmarek, M. R. Lewis, R. Rottgers, A. Sciandra, M. Stramska, M. S. Twardowski, B. A. Franz, and H. Claustre (2008), Relationships between the surface concentration of particulate organic carbon and optical properties in the eastern South Pacific and eastern Atlantic Oceans, *Biogeosciences*, *5*, 171–201.
- Stumpf, R. P. (1988), Sediment transport in Chesapeake Bay during floods; analysis using satellite and surface observations, *Journal of Coastal Research*, *4*, 1–15.
- Stumpf, R. P., and J. R. Pennock (1989), Calibration of a general optical equation for remote sensing of suspended sediments in a moderately turbid estuary, *J. Geophys. Res.*, *94*(C10), 14363–14371.
- Trefry, J. H., S. Metz, T. A. Nelsen, R. P. Trocine, and B. J. Eadie (1994), Transport of particulate organic-carbon by the mississippi river and its fate in the Gulf-of-Mexico, *Estuaries*, *17*, 839–849.
- Walker, N. D., and N. N. Rabalais (2006), Relationships among Satellite Chlorophyll a, River Inputs, and Hypoxia on the Louisiana Continental Shelf, Gulf of Mexico, *Estuaries and Coasts*, *29*(6B), 1081–1093.

Wang, X.-C., R. F. Chen, and G. B. Gardner (2004), Sources and transport of dissolved and particulate organic carbon in the Mississippi River estuary and adjacent coastal waters of the northern Gulf of Mexico, *Marine Chemistry*, 89 241- 256.